



TITLE:

Nonprincipal Block of $SL(2, q)$ (Cohomology Theory of Finite Groups and Related Topics)

AUTHOR(S):

Yoshii, Yutaka

CITATION:

Yoshii, Yutaka. Nonprincipal Block of $SL(2, q)$ (Cohomology Theory of Finite Groups and Related Topics). 数理解析研究所講究録 2008, 1581: 121-125

ISSUE DATE:

2008-02

URL:

<http://hdl.handle.net/2433/81423>

RIGHT:

Nonprincipal Block of $SL(2, q)$

Yutaka Yoshii (吉井 豊)

Division of Mathematical Science and Physics, Chiba Univ. (千葉大学自然科学研究科)

Abstract

We shall claim that Broué's abelian defect group conjecture holds for the nonprincipal p -block of $SL(2, p^n)$.

1 Introduction

Let G be a finite group and P a p -subgroup of G . The next theorem is one of the most important theorems on the block theory of finite groups:

Brauer's First Main Theorem. *There is one to one correspondence between the blocks of kG with defect group P and the blocks of $kN_G(P)$ with defect group P .*

The correspondence is called *Brauer correspondence*. The following conjecture is our main problem:

Broué's Abelian Defect Group Conjecture. *Suppose that A is a block of kG with an abelian defect group P and that B is the Brauer correspondent of A (in $N_G(P)$). Then is A derived equivalent to B ?*

If $G = SL(2, q)$ where $q = p^n$, it has been proved that the conjecture is true for the principal block by T.Okuyama (see [6]). Even in the nonprincipal case, the conjecture was proved to be true for $n = 2$ by M.Holloway (see [4]), but it has not been known if the conjecture is true for $n \geq 3$ yet. However, it has turned out that it can be proved to be true even for $n \geq 3$ by imitating Okuyama's proof [6].

The Main Result. *If $G = SL(2, q)$ where $q = p^n$, Broué's abelian defect group conjecture is true for the nonprincipal block of kG .*

We shall explain about derived equivalences. Let k be an algebraically closed field of characteristic $p > 0$, let A and B be finite dimensional k -algebras, $\text{mod-}A$ the category consisting of all finite dimensional right A -modules, $\text{proj-}A$ the full subcategory of $\text{mod-}A$ consisting of all finite dimensional right projective A -modules, $K^b(\text{mod-}A)$ the homotopy category consisting of all bounded complexes of finite dimensional right A -modules, and $K^b(\text{proj-}A)$ the homotopy category consisting of all bounded complexes of finite dimensional right projective A -modules. We say that A is *derived equivalent* to B if $K^b(\text{proj-}A)$ is equivalent to $K^b(\text{proj-}B)$ as triangulated categories. The next theorem is a criterion for derived equivalence:

Theorem(Rickard [7]). *The following are equivalent.*

- (a) A is derived equivalent to B .
- (b) There is a complex $T^\bullet \in K^b(\text{proj-}A)$ with $B \cong \text{End}_{K^b(\text{proj-}A)}(T^\bullet)$ such that
 - (i) $\text{Hom}_{K^b(\text{proj-}A)}(T^\bullet, T^\bullet[i]) = 0$ for any $i \neq 0$.
 - (ii) If $\text{add}(T^\bullet)$ is the full subcategory of $K^b(\text{proj-}A)$ consisting of all direct summands of all direct sums of T^\bullet , then it generates the triangulated category $K^b(\text{proj-}A)$.

We call T^\bullet a *tilting complex* for A .

2 $SL(2, q)$

Set $G = SL(2, q)$ where $q = p^n$. In this section, we shall state some facts of representations of kG . Set

$$P = \left\{ \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \middle| b \in \mathbb{F}_q \right\},$$

$$D = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \middle| a \in \mathbb{F}_q^\times \right\},$$

and

$$H = N_G(P) = \left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} \mid a \in \mathbb{F}_q^\times, b \in \mathbb{F}_q \right\},$$

where P is a Sylow p -subgroup of G and hence is isomorphic to the elementary abelian group $C_p \times \cdots \times C_p$ (n times), D is isomorphic to C_{q-1} , and H is the semidirect product $P \rtimes D$.

Considering a nonprincipal block, we assume $p \neq 2$ in the rest of the article (if $p = 2$, kG has no nonprincipal blocks with full defect). Now we have the block decompositions $kG = A_0 \oplus A_1 \oplus A_2$, where A_0 is the principal block, A_1 is a nonprincipal block with full defect, and A_2 has defect zero, and $kN_G(P) = B_0 \oplus B_1$, where B_0 and B_1 are the Brauer correspondents of A_0 and A_1 respectively. It is well known that all nonisomorphic simple kG -modules are indexed by $\{0, 1, 2, \dots, q-1\}$, where $\{0, 2, \dots, q-3\}$, $\{1, 3, \dots, q-2\}$ and $\{q-1\}$ correspond to A_0 , A_1 and A_2 respectively; and all nonisomorphic simple $kN_G(P)$ -modules are indexed by $\{0, 1, 2, \dots, q-2\}$, where $\{0, 2, \dots, q-3\}$ and $\{1, 3, \dots, q-2\}$ correspond to B_0 and B_1 respectively (see [3] or [6]).

3 Outline of Proof

Set $\Lambda = \{0, 1, 2, \dots, q-1\}$, $I = I_{\text{odd}} = \{1, 3, 5, \dots, q-2\}$. For $\lambda \in \Lambda - \{q-1\}$, set

$$\tilde{\lambda} = \begin{cases} 0 & (\text{if } \lambda = 0) \\ q-1-\lambda & (\text{if } \lambda \neq 0), \end{cases}$$

and for a subset $\Omega \subseteq \Lambda - \{q-1\}$, set $\tilde{\Omega} = \{\tilde{\lambda} \mid \lambda \in \Omega\}$. Then for any simple $kN_G(P)$ -module, $T_{\tilde{\lambda}}$ is isomorphic to the dual module T_{λ}^* of T_{λ} , and note that " \sim " is a permutation on $\Lambda - \{q-1\}$ of order 2. Moreover, we define an equivalence relation " \sim " on $\Lambda - \{q-1\}$ by

$$\lambda \sim \mu \stackrel{\text{def}}{\iff} \text{There exists some } j \in \{0, 1, \dots, n-1\} \text{ such that } \lambda \equiv p^j \mu \pmod{q-1}.$$

Note that I is closed under the equivalence relation.

We define equivalence classes (with respect to " \sim ") $J_{-1}, J_0, J_1, \dots, J_s$ as follows (cf. Okuyama [6, §2]):

Let J_{-1}, \tilde{J}_{-1} be empty sets (by convention), J_0 the class containing 1, and J_i the class containing the smallest $\lambda_i \notin \bigcup_{u=-1}^{i-1} (J_u \cup \tilde{J}_u)$ for $i \geq 1$. We repeat this procedure until s satisfies $I = \bigcup_{u=-1}^s (J_u \cup \tilde{J}_u)$.

Now we can construct derived equivalent k -algebras $A^0, A^1, \dots, A^s, A^{s+1}$ as follows (cf. Okuyama [6, §3]):

First, set $A^0 = A$. Then for $1 \leq t \leq s+1$, we define A^t as an endomorphism algebra of a tilting complex for A^{t-1} determined by J_{t-1} which is seen in [6, §1].

Then, we can show that A^{s+1} is isomorphic to B as k -algebras like Okuyama [6, §3], so we obtain the main result.

References

- [1] H.H.Andersen, J.Jørgensen, and P.Landrock, *The projective indecomposable modules of $SL(2, p^n)$* , Proc. London Math. Soc.(3) **46** (1983), no.1, 38-52.
- [2] M.Broué, *Isométries parfaites, types de blocs, catégories dérivées*, Astérisque **181-182** (1990), 61-92.
- [3] P.W.A.M. van Ham, T.A.Springer, and van der Wel, *On the Cartan invariant of $SL(2, \mathbb{F}_q)$* , Comm. Alg. **10(14)** (1982), 1565-1588.
- [4] M.Holloway, *Derived equivalences for group algebras*, Ph.D Thesis, University of Bristol (2001).
- [5] T.Okuyama, *Some examples of derived equivalent blocks of finite groups*, preprint (1998).
- [6] T.Okuyama, *Derived equivalence in $SL(2, q)$* , preprint (2000).
- [7] J.Rickard, *Morita theory for derived categories*, J. London Math. Soc. (2) **39** (1989), 436-456.
- [8] J.Rickard, *Derived equivalences as derived functors*, J. London Math. Soc. (2) **43** (1991), 37-48.
- [9] J.Rickard, *Splendid equivalences: Derived categories and permutation modules*, Proc. London Math. Soc.(3) **72** (1996), 331-358.
- [10] R.Rouquier, *From stable equivalences to Rickard equivalences for blocks with cyclic defect*, Groups '93 Galway/St Andrews II (C.M.Campbell

et al., eds.), vol. 212, London Math. Soc. Lecture Note Series (1995), pp.512-523.

- [11] R.Rouquier, *The derived category of blocks with cyclic defect groups*, Derived Equivalences for Group Rings (S.König and A.Zimmermann), vol. 1685, Springer Lecture Notes in Mathematics (1998), pp.199-220.
- [12] R.Rouquier, *Block theory via stable and Rickard equivalences*, Modular representation theory of finite groups (eds. M.J.Collins, B.J.Parshall and L.L.Scott), pp.101-146, de Gruyter, Berlin (2001).